



Strangeness Report

Federico Antinori
INFN Padova - Italy



Contents

→ I concentrate on strange particle abundances (yields and ratios)

- QGP predictions
- Enhancement pattern
- Hadronic transport
- Thermal models
- Beyond equilibrium?
- The Φ
- Extras
- Conclusions

not covered: Resonances (Fachini), Spectra (Velkovska), Flow (Retiere),
Pentaquarks (Jaffe), HBT (Magestro)

Two historic QGP predictions

- **restoration of χ symmetry \rightarrow increased production of s**


- mass of strange quark in QGP expected to go back to current value

- $m_s \sim 150 \text{ MeV} \sim T_c$

\rightarrow copious production of $s\bar{s}$ pairs, mostly by gg fusion

[Rafelski: Phys. Rep. 88 (1982) 331]

[Rafelski-Müller: P. R. Lett. 48 (1982) 1066]



- **deconfinement \rightarrow stronger effect for multi-strange**


- can be built using uncorrelated s quarks produced in independent microscopic reactions

\rightarrow **strangeness enhancement increasing with strangeness content**

[Koch, Müller & Rafelski: Phys. Rep. 142 (1986) 167]


and indeed...

- @ top SPS (NA57, $\sqrt{s_{NN}} = 17.3 \text{ GeV}$)




and indeed...

- and (yesterday's news) @ RHIC (STAR, $\sqrt{s_{NN}} = 200$ GeV)






and indeed...

- and (yesterday's news) @ low E SPS (NA57, $\sqrt{s_{NN}} = 8.8$ GeV)



and indeed...



- qualitatively similar picture emerging from 8.8 to 200 GeV

Alternative definition of pA enhancement?

e.g.: [Fischer: Nucl. Phys. A 715 (2003) 118]

- two-component model of particle production:

$$\begin{aligned} \rightarrow \text{in pA: } Y_{pA} &= [\nu/2 + 1/2] Y_{pp} & (\nu = \text{number of collisions}) \\ &= N_{\text{part}} Y_{pp}/2 \end{aligned}$$

- enhancement "blamed" on projectile component only:

usual enhancement: E

alternative enhancement: F

$$(Y/N_{\text{part}})_{pA} = E \cdot (Y/N_{\text{part}})_{pp}$$

$$\rightarrow Y_{pA} = E [\nu/2 + 1/2] Y_{pp}$$

$$Y_{pA} = [\nu/2 + F/2] Y_{pp}$$

reasonable for baryon number transfer,
but do e.g. strange quarks know if they
come from projectile or target ?

- then: $(F - 1) = (E - 1) N_{\text{part}}$

\rightarrow enhancement within pA appears even if pA yields $\propto N_{\text{part}}$

Hadronic transport

- Hadronic transport codes:
 - do reasonably well on singly strange particles
 - but fail to reproduce the production of multi-strange particles at SPS and RHIC
 - see for instance:
 - [Soff et al.: Phys. Lett. B471 (1999) 89],
 - [C.Greiner: nucl-th/0208080 and references there],
 - [STAR: nucl-ex/0307024],
 - [Huovinen & Kapusta: nucl-th/0310051]

[Huovinen & Kapusta: nucl-th/0310051]

Hadronic transport

- Hadronic transport codes:
 - do reasonably well on singly strange particles
 - but fail to reproduce the production of multi-strange particles at SPS and RHIC

● see for example

- [Schnitzler et al., 1999]
- [Cassorina et al., 1999]
- [Schnitzler et al., 1999]
- [Huovinen & Kapusta, 2003]





FIG. 1. The ratio of the calculated abundance of the indicated species to the chemical equilibrium value as a function of the local temperature. The upper set of curves start with the baryons in equilibrium at T_c , the lower set start with no baryons.

[Huovinen & Kapusta: nucl-th/0310051]

Hadronic transport



- Hadronic transport codes:
 - do reasonably well on singly strange particles
 - but fail to reproduce the production of multi-strange particles at SPS and RHIC
 - see for instance:
 - [Soff et al.: Phys. Lett. B471 (1999) 89],
 - [C.Greiner: nucl-th/0208080 and references there],
 - [STAR: nucl-ex/0307024],
 - [Huovinen & Kapusta: nucl-th/0310051]
- they get closer if:
 - masses are reduced towards chiral values
 - or string tension is enhanced with respect to ee/pp/pA
 - enhanced contribution from inelastic scattering during expansion
- Activity is still ongoing:
 - e.g.: [Capella: nucl-th/0303045; Subrata Pal et al.: nucl-th/0106074]
 - hadronic models are getting more and more exotic...



[Huovinen & Kapusta: nucl-th/0310051]

Meanwhile at the AGS...

- Nice new measurement: Ξ^- production close to threshold, Au-Au @ 6 A GeV (E895) [P.Chung et al.: nucl-ex/0302021]
- At this energy RQMD works fine!




Thermal fits

- Thermal fits don't do too bad!

e.g.: brand new @ RHIC 200


- $T_{ch} = 160 \pm 10 \text{ MeV}$
- $\mu_B = 24 \pm 5 \text{ MeV}$

(STAR Preliminary)




- relative particle abundances ~ as expected at thermodynamical equilibrium **for grand-canonical system**, even for rare, multi-strange particles

T systematics



- it looks like Hagedorn was right!
 - if the resonance mass spectrum grows exponentially (and this seems to be the case), there is a maximum possible temperature for a system of hadrons
 - indeed, we don't seem to be able to get a system of hadrons with a temperature beyond $T_{\max} \sim 170$ MeV!


T vs μ_B systematics



- the extracted freeze-out points at SPS and RHIC lay very close to the predicted QGP phase boundary

Canonical vs Grand Canonical



- Energy penalty to create a strange particle:
 - Canonical:
computed taking into account also energy to create companion to ensure conservation of strangeness
 - Grand Canonical limit:
just due to creation of particle itself. The rest of the system acts as a reservoir and "picks up the slack"





- removal of canonical suppression
 - increases with strangeness
 - \sim observed enhancements
 - detailed centrality dependence not reproduced (very crude modelling)

[Hamieh et al.: Phys. Lett. B486 (2000) 61]

● @ RHIC:



- @ RHIC:



- decrease of enhancements from SPS predicted
- order of magnitude of enhancements close
- again, model for V_{corr} vs N_{part} seems to be inadequate

Does this *explain* the observed enhancement pattern?

- a system in eq., if it is large enough, is in GC eq., but being large in itself is not a sufficient condition for being GC!
 - if AA colls. were just a superposition of pp, they would have to be treated canonically all the same!
- the system must also **know** it is large...
 - it must **know** that an $\bar{\Omega}^+$ generated here can be compensated by, say, an Ω^- on the other side of the fireball!
 - what counts is the *correlation volume*
- "Canonical Suppression is removed!"
 - an *observation*, not an *explanation*


How is Canonical Suppression removed?

how does the system know it is large?

how can information travel so quickly through the system?

→ not by conventional hadronic transport (no time!)

- natural if the system is coming back from deconfinement



How is Canonical Suppression removed?

how does the system know it is large?

how can information travel so quickly through the system?

→ not by conventional hadronic transport (no time!)

- natural if the system is coming back from deconfinement
- two recent ideas:
 - quantum coherence in the fireball [Stock: hep-ph/0312039]
 - can it be developed into testable model?
 - what about pA?
 - multi-particle interactions [Braun-Munzinger et al.: nucl-th/0311005]
 - only important close to phase boundary
 - if true, we are actually *measuring* T_c
 - microscopic mechanism to restore abundances to hadronic eq.

An equilibrium *pre*-diction


- mid-rapidity multi-strange/ Λ @ 40 GeV SPS

NA57 vs [Becattini: private communication]


- $\gamma_s = 1$, T from 4π fit to NA49 40 GeV [Becattini et al.: hep-ph/0310049]
 - using π^\pm , K^\pm , ϕ , Λ and $\bar{\Lambda}$
- Λ and $\bar{\Lambda}$ dN/dy from NA57 fix μ_B and normalization
- Prediction of Ξ^- , Ξ^+ , Ω^- and $\bar{\Omega}^+$ dN/dy
 - total $\chi^2/\text{dof} = 1.4$ against values measured by NA57

→ No parameter for multi-strange
again, not bad...

Physics beyond equilibrium?
→ look at fine details



$\Lambda(1520)$



- $\Lambda(1520)$ deviates from systematics ($\sim 1/2$)

e.g.: [Markert: J. Phys. G 28 (2002) 1753]


→ we look for in-medium effects

we take fits very seriously!

Fit to NA49 data
 [Becattini et al.: hep-ph/0310049]

Resonances @ RHIC

- new: $\Lambda(1520)$, $K^*(892)$, $\Delta(1232)$ in STAR at 200 GeV



- resonances show larger deviations from statistical fit
- rescattering vs regeneration?
 - see talk by P. Fachini

Non equilibrium: γ_s

- 4π yield ratios sometimes preferred for thermal fits
 - to be safe in case strangeness ends up too far away in rapidity from where originally produced
 - (but can one assume global equilibrium with same T and same μ_B at different rapidities?)
- in this case $\gamma_s \sim 0.7 - 0.8$ must be introduced
 - strangeness undersaturation factor:
for each particle, a factor $\gamma_s^{N(s+\bar{s})}$
e.g.: [Becattini et al.: hep-ph/0310049]
 - could it also mimic residual canonical suppression?

Non equilibrium: γ_s & γ_q

see e.g.: [Rafelski & Letessier: hep-ph/0309030]

- γ_q controls overall abundance of $q\bar{q}$
- old γ_s becomes γ_s/γ_q
- w.r.t. previous, additional factor γ_q^B for baryons
- fits get beautiful e.g.: @ RHIC \rightarrow
[see also Becattini et al.: hep-ph/0310049]
(but additional parameter)

- @ RHIC 130:


$$\gamma_q = 1.6 \pm 0.2 \quad \gamma_s/\gamma_q = 1.3 \pm 0.1$$

- strangeness actually enhanced above hadronic equilibrium values?

ratio	RHIC-130	non-eq fit	χ^2
π^+/p	9.5 ± 1.4	9.07	1.15
π^-/p	13.4 ± 0.9	13.15	0.08
p/h^-	—	0.0459	—
Λ_c/h^-	0.059 ± 0.004	0.0509	4.11
$\bar{\Lambda}_c/h^-$	0.042 ± 0.004	0.0379	1.04
Ξ_c^-/h^-	0.0079 ± 0.0012	0.00805	0.01
$\bar{\Xi}_c^-/h^-$	0.0066 ± 0.001	0.00645	0.02
Ω/h^-	$(12 \pm 5)10^{-4}$	$13.2 \cdot 10^{-4}$	0.06
$(\bar{\Omega} + \Omega)/h^-$	$(22 \pm 6.5)10^{-4}$	$24.8 \cdot 10^{-4}$	0.19
Λ_c/p	0.90 ± 0.12	0.747	1.63
$\bar{\Lambda}_c/p$	0.93 ± 0.19	0.826	0.30
Ξ^-/Λ	0.193 ± 0.03	0.189	0.02
$\bar{\Xi}^-/\bar{\Lambda}$	0.219 ± 0.035	0.207	0.12
Ω/Ξ^-	—	0.164	—
$\bar{\Omega}/\bar{\Xi}^-$	—	0.180	—
$(\bar{\Omega} + \Omega)/(\bar{\Xi} + \Xi)$	0.150 ± 0.04	0.171	0.28
p/p	0.71 ± 0.06	0.674	0.36
Λ_c/Λ_c	0.71 ± 0.04	0.745	0.78
Ξ/Ξ	0.83 ± 0.08	0.801	0.13
$\bar{\Omega}/\Omega$	0.95 ± 0.1	0.878	0.51
K^+/π^+	0.17 ± 0.02	0.195	1.59
K^-/π^-	0.15 ± 0.02	0.180	2.28
K^-/K^+	0.87 ± 0.07	0.923	0.57
K^{*0}/K^-	0.26 ± 0.08	0.231	0.13
ϕ/h^-	0.02 ± 0.002	0.0212	0.37
ϕ/K^-	0.15 ± 0.03	0.148	0.00
ϕ/K^{*0}	0.595 ± 0.24	0.639	0.03


$\sqrt{s_{NN}}$ [GeV]	200	130	200	130
T [MeV]	143 ± 7	144 ± 3	160 ± 8	160 ± 4
μ_b [MeV]	21.5 ± 31	29.2 ± 4.5	24.5 ± 3	31.4 ± 4.5
μ_s [MeV]	2.5 ± 0.2	3.1 ± 0.2	2.9 ± 0.2	3.6 ± 0.2
$\mu_{\bar{s}}$ [MeV]	4.7 ± 0.4	6.6 ± 0.4	5.3 ± 0.4	6.9 ± 0.4
γ_q	$1.6 \pm 0.3^*$	$1.6 \pm 0.2^*$	1^*	1^*
γ_s/γ_q	1.2 ± 0.15	1.3 ± 0.1	1.0 ± 0.1	1.13 ± 0.06
χ^2/dof	2.9/6	15.8/24	4.5/7	32.2/25
P_{true}	90%+	95%+	65%	15%

- pentaquark production (if statistical) would be particularly sensitive to need for γ_q !
 - additional γ_q^2 factor for Θ^+ w.r.t. other models
e.g.: [Letessier et al.: hep-ph/0310188, Becattini et al.: hep-ph/0310049]



Hyperon enhancements at 40 GeV

- enhancements vs N_{wound} steeper at 40 than at 160
- C to GC predicts a factor 4 - 5 larger Ξ^- enhancement at 40 GeV than at 160, not seen in the data




[Bruno (NA57): yesterday]

[Tounsi & Redlich: hep-ph/0111159]

- perhaps yields trying to shoot up but don't go all the way?
 - on the other hand, equilibrium fit not too bad... [Becattini: priv. comm]


Marek's Horn

- Change of behaviour of K^+/π^+ in A-A around 30 GeV
 - K^-/π^- has a smooth behaviour with energy




Brahms: K^\pm vs y

- caution! K^\pm sensitive to baryon density...




[Brahms: Phys. Rev. Lett. 90 (2003) 102301]




[D. Ouerdane, PhD Dissertation, Copenhagen University, Aug. 03]

- ... and $\lambda_s [= 2\langle s\bar{s} \rangle / (\langle u\bar{u} \rangle + \langle d\bar{d} \rangle)]$ max. around 30 GeV:



[Braun-Munzinger et al.: NPA697(2002)902]


- so one expects a broad maximum in K^+/π^+ about there...



- but nothing so sharp!
→ it looks like we have some anomaly...

Absolute values of hyperon dN/dy


see talk by D.Elia, Friday, P2




- absolute values of $\bar{\Lambda}$ and $\bar{\Xi}^+$ dN/dy go up with \sqrt{s} ...
- absolute values of Λ and Ξ^- dN/dy \sim constant from $\sqrt{s_{NN}} = 8.8$ to 130 GeV !
- Statistical fits: T and μ_B only sensitive to particle ratios
 - overall normalization of yields controlled by fitted V parameter
- T , μ_B and V all vary with energy, but in such a way as to ensure Λ , Ξ^- yields stays constant ...
 - there must be a simple reason ...

The mysterious Φ

- Φ fundamental to understand γ_s vs C suppression
 - γ_s^2 , but no C suppression
- the NA49 - NA50 discrepancy is still with us...
- NA49: small deviation from PDG mass seen in all three Pb+Pb data sets, but not in pp



→ NA60 should provide
high stats, low p_T $\Phi \rightarrow \mu\mu$





see [NA49 (Friese): SQM 03]

First Φ signal from NA60


- peripheral In-In
- expected $> 100k \Phi \rightarrow \mu^+\mu^-$
- expected mass resolution: $\sim 20 \text{ MeV}$
- also looking at $\Phi \rightarrow K^+K^-$

ω ϕ



Φenix


- $\Phi \rightarrow K^+K^-$ (min bias 200 GeV)



Mass (GeV/c²)

- Mass and width agree within errors with PDG values

- $\Phi \rightarrow e^+e^-$ (min bias 200 GeV)



- Preliminary yields:

$$\phi \rightarrow e^+e^- : \quad \frac{dN}{dy} = 5.4 \pm 2.5 (stat) {}^{+3.4}_{-2.8} (sys)$$

$$\phi \rightarrow K^+K^- : \quad \frac{dN}{dy} = 2.01 \pm 0.22 (stat) {}^{+1.01}_{-0.52} (sys)$$


see [Phenix (Maguire): SQM 03]

Extras...


some results on spectra that I could not
resist showing...

Very low p_T π , K , p from Phobos !

- tracks characterized by range and energy loss in Si layers
 - particle id.
 - momentum within 5%
- unique constraints for transport, hydro, thermal models




Rcp for multistrange !




- Φ seems to follow K_S^0 more than Λ
 → meson-baryon, not mass?

v2 for multistrange !




scaled with $n(\text{quarks})$

- scaling seems to work (not for pions...)
→ parton recombination?




m_T scaling in pp...





not absolute...

...but if one rescales:



... but not in Au-Au



data



← p-p →

← Au-Au →



data / power law



Conclusions

- Evidence for high degree of chemical equilibration
 - even for rare, multistrange particles
 - QGP prediction !
 - freeze-out curve traces expected phase boundary at SPS - RHIC
 - hadronic transport does not seem to be able to do it (no time!)
- RHIC opening up exciting new windows for strangeness !
 - elliptic flow, high p_T , high statistics reference data (pp, dAu)
- Strangeness perhaps enhanced above hadronic equilibrium !
 - stand-alone smoking gun?

- we must understand (non-)equilibrium
 - we need highest possible precision data on yields and ratios (RHIC!)
 - (we need theory to provide agreed statistical approach)
 - we need to understand the Φ
 - precision data from NA60 and Phenix
- we must understand what's going on in the region
 - $\sqrt{s_{NN}} = 5 - 10 \text{ GeV}$
 - low E SPS ?
 - SIS 200 @ GSI !



Credits

- Many thanks for discussions and material to:
 - Francesco Becattini
 - Rene Bellwied
 - Giuseppe Bruno
 - Helen Caines
 - Domenico Elia
 - Marek Gazdzicki
 - Adam Jacholkowski
 - Carlos Lourenço
 - Djamel Ouerdane
 - Emanuele Quercigh
 - Jan Rafelski
 - Karel Šafařík
 - Reinhard Stock
 - Julia Velkovska